arXiv:1303.2263v1 [math.CO] 9 Mar 2013

Fan-type degree condition restricted to triples of induced subgraphs ensuring Hamiltonicity[*]

Bo Ning[†]

Department of Applied Mathematics, School of Science,

Northwestern Polytechnical University, Xi'an, Shaanxi 710072, P.R. China

Abstract

In 1984, Fan gave a sufficient condition for Hamilton cycles involving maximum degree of every pair of vertices at distance two. Motivated by Fan's result, we say that an induced subgraph H of G is f-heavy if for every pair of vertices $u, v \in V(H)$, $d_H(u, v) = 2$ implies that $\max\{d(u), d(v)\} \geq n/2$. For a given graph R, G is called R-f-heavy if every induced subgraph of G isomorphic to R is f-heavy. For a family $\mathcal{R}$ of graphs, G is called $\mathcal{R}$-f-heavy if G is R-f-heavy for every $R \in \mathcal{R}$. In this note we show that every 2-connected graph G has a Hamilton cycle if G is $\{K_{1,3}, P_7, D\}$-f-heavy or $\{K_{1,3}, P_7, H\}$-f-heavy, where P_7 is a path of length 6, H consists of two triangles sharing a common vertex, and D is obtained by attaching two disjoint paths of length 2 to two vertices of a triangle. Our result extends a previous theorem of Broersma, Ryjáček and Schiermeyer [H.J. Broersma, Z. Ryjáček, I. Schiermeyer, Dirac's minimum degree condition restricted to claws, Discrete Math. 167-168 (1997) 155-166] on Hamiltonicity of 2-connected graphs.

Keywords: Hamilton cycle; Fan-type degree condition; Induced subgraph; Claw

AMS Subject Classification (2000): 05C38 05C45

1 Introduction

We use Bondy and Murty [3] for terminology and notation not defined here and consider simple graphs only.

---

[*] Address for correspondence: Department of Applied Mathematics, Northwestern Polytechnical University, Xi'an, P.R.China.

[†] E-mail address: ningbo_math84@mail.nwpu.edu.cn.

Let G be a graph on n vertices. If a subgraph G' of G contains every edge $xy \in E(G)$ with $x, y \in V(G')$, then G' is called an *induced subgraph* of G. For a given graph R, G is *R-free* if G contains no induced subgraph isomorphic to R, and *R-f-heavy* if for every induced subgraph G' of G isomorphic to R and every pair of vertices $u, v \in V(G')$, $d_{G'}(u,v) = 2$ implies that $max\{d(u), d(v)\} \geq n/2$. For a family $\mathcal{R}$ of graphs, G is *$\mathcal{R}$-free* (*$\mathcal{R}$-f-heavy*) if G is *R-free* (R-f-heavy) for every $R \in \mathcal{R}$. Note that every $\mathcal{R}$-free graph is also $\mathcal{R}$-f-heavy.

The graph $K_{1,3}$ is called a *claw* and its vertex of degree one is called an *end-vertex*. Following [4], a graph G is 2-*heavy* if at least two end-vertices of each induced claw of G have degree at least $n/2$. Throughout this note, we use *claw-free* (2-heavy) instead of $K_{1,3}$-free ($K_{1,3}$-f-heavy).

A graph is called *Hamiltonian* if it contains a spanning cycle. In 1984, Fan [8] gave a sufficient condition for the existence of Hamilton cycles in 2-connected graphs as follows.

Theorem 1 (Fan [8]). *Let G be a 2-connected graph on $n \geq 3$ vertices. If $\max\{d(x), d(y)\} \geq n/2$ for every pair of vertices x and y with $d(x,y) = 2$, then G is Hamiltonian.*

Obviously, Fan's result is equivalent to every 2-connected P_3-f-heavy graph is Hamiltonian. Thus we can naturally pose this problem: which two connected graphs R and S other than P_3 imply that every 2-connected $\{R, S\}$-f-heavy graph is Hamiltonian? In fact, Bedrossian, Chen and Schelp [1], Chen, Wei and Zhang [6, 7], and Li, Wei and Gao [11] got some results related to this problem, respectively. Recently, this problem has been completely solved in [12].

The main purpose of this note is to find triples of Fan-type heavy subgraphs which ensure Hamiltonicity of graphs. First, there are a series of results about Hamilton cycles based on triples of forbidden subgraphs. The following two results are well-known ones in this direction.

Theorem 2 (Broersma and Veldman [5]). *Let G be a 2-connected graph. If G is claw-free and $\{P_7, D\}$-free, then G is Hamiltonian. (see Fig. 1)*

Theorem 3 (Faudree, Ryjáček and Schiermeyer [9]). *Let G be a 2-connected graph. If G is claw-free and $\{P_7, H\}$-free, then G is Hamiltonian. (see Fig. 1)*



D (Deer) H (Hourglass)

Fig. 1. Graphs D and H.

Broersma et al. [4] extended Theorems 2 and 3 to a larger class of 2-heavy graphs.

Theorem 4 (Broersma, Ryjáček and Schiermeyer [4])**.** *Let G be a 2-connected graph. If G is 2-heavy, and moreover $\{P_7, D\}$-free or $\{P_7, H\}$-free, then G is Hamiltonian.*

By restricting Fan's condition to triples of induced subgraphs of graphs, we extends Theorem 4 as follows.

Theorem 5. *Let G be a 2-connected graph. If G is $\{K_{1,3}, P_7, D\}$-f-heavy or $\{K_{1,3}, P_7, H\}$-f-heavy, then G is Hamiltonian.*

Remark 1. The graph provided in the following shows that Theorem 5 indeed strengthen Theorem 4. Let $n \geq 16$ be an even integer and $K_{n/2} + K_{n/2-7}$ denotes the union of two complete graphs $K_{n/2}$ and $K_{n/2-7}$. We construct the graph G with $V(G) = V(K_{n/2} + K_{n/2-7}) \cup \{x, y, z, u, v, w, t\}$ and $E(G) = E(K_{n/2} + K_{n/2-7}) \cup \{xy, xz, yz, yw, wu, zt, tv\} \cup \{xx', yx', zx' : x' \in V(K_{n/2})\} \cup \{uy', vy' : y' \in V(K_{n/2-7})\}$. It is easy to see that G is a Hamiltonian graph satisfying the condition of Theorem 5, but not the condition of Theorem 1 or 4.

The rest of this note is organized as follows. In Section 2, we will give additional terminology and notation and list some useful lemmas. The proof of Theorem 5 will be postponed to Section 3.

2 Preliminaries

Let G be a graph and H be a path or a cycle of G. We denote by $\overrightarrow{H}$ the path (or the cycle) with a given orientation, and by $\overleftarrow{H}$ the same subgraph with the reverse orientation. For two vertices $x, y \in V(H)$, $\overrightarrow{H}[x, y]$ is denoted by the consecutive vertices from x to y

in H by the direction specified by $\overrightarrow{H}$, and $\overleftarrow{H}[y,x]$ is the same vertices with the reverse order. For a vertex $x \in V(H)$, x^+ denotes the *successor* of x on $\overrightarrow{H}$, and x^- denotes its *predecessor*.

Recall that a vertex of a graph G on n vertices is *heavy* if its degree is at least $n/2$; otherwise, it is *light*. A cycle C of G is called a *heavy cycle* if it contains all the heavy vertices of G. Following [10], we use $\widetilde{E}(G)$ to denote the set $\{xy : xy \in E(G) \text{ or } d(x) + d(y) \geq n, x, y \in V(G)\}$. Let $k \geq 3$ be an integer. A sequence of vertices $C = v_1v_2 \ldots v_kv_1$ is called an *Ore-cycle* or briefly, *o-cycle* of G, if for every $i \in \{1, \cdots, k\}$, there holds $v_iv_{i+1} \in \widetilde{E}(G)$, where the indices are taken modulo k.

Lemma 1 (Bollobás and Brightwell [2], Shi [13])**.** *Let G be a 2-connected graph. Then G contains a heavy cycle.*

Lemma 2 (Li, Ryjáček, Wang and Zhang [10])**.** *Let G be a graph and C be an o-cycle of G. Then there exists a cycle C' of G such that $V(C) \subseteq V(C')$.*

3 Proof of Theorem 5

By Lemma 1, G contains a heavy cycle. Let C be a longest heavy cycle of G, fixed an orientation. Suppose that G is not Hamiltonian. Since G is 2-connected, there is a path of length at least 2, internally-disjoint with C, that connects two vertices of C. Let $\overrightarrow{P} = w_0w_1 \ldots w_rw_{r+1}$ be such a path with r as small as possible, where $w_0 = u \in V(C)$ and $w_{r+1} = v \in V(C)$.

Claim 1. Let $x \in V(P) \backslash \{u, v\}$ and $y \in \{u^-, u^+, v^-, v^+\}$. Then $xy \notin \widetilde{E}(G)$.

Proof. Without loss of generality, assume that $y = u^-$. Suppose that $xy \in \widetilde{E}(G)$. Then $C' = yx\overleftarrow{P}[x,u]\overrightarrow{C}[u,y]$ is an o-cycle containing all the vertices of C and longer than C. By Lemma 2, there is a longer cycle containing all the vertices in C, that is, a longer heavy cycle in G, a contradiction. The other assertions can be proved similarly. □

Claim 2. $u^-u^+ \in \widetilde{E}(G)$, $v^-v^+ \in \widetilde{E}(G)$.

Proof. Suppose that $u^-u^+ \notin E(G)$. By Claim 1, $\{u, u^-, u^+, w_1\}$ induces a claw. By the choice of C, w_1 is light. Since G is 2-heavy, we have $d(u^-) + d(u^+) \geq n$. This implies that $u^-u^+ \in \widetilde{E}(G)$. Similarly, we can prove the other assertion. □

Claim 3. $uv^{\pm} \notin \widetilde{E}(G)$, $vu^{\pm} \notin \widetilde{E}(G)$, $u^-v^- \notin \widetilde{E}(G)$, $u^+v^+ \notin \widetilde{E}(G)$.

Proof. Suppose that $uv^- \in \widetilde{E}(G)$. By Claim 2, $u^+u^- \in \widetilde{E}(G)$. Then $C' = uv^-\overleftarrow{C}[v^-, u^+]u^+u^-\overleftarrow{C}[u^-, v]\overleftarrow{P}[v, u]$ is an o-cycle containing all vertices in C and longer than C, a contradiction. Suppose $u^-v^- \in \widetilde{E}(G)$. Then $C' = u^-v^-\overleftarrow{C}[v^-, u]\overrightarrow{P}\overrightarrow{C}[v, u^-]$ is an o-cycle containing all vertices in C and longer than C, a contradiction. The other assertions can be proved similarly. □

Let y_1 be the first vertex on $\overrightarrow{C}[u, v]$ such that $uy_1 \notin E(G)$, y_2 be the first vertex on $\overrightarrow{C}[v, u]$ such that $vy_2 \notin E(G)$. By Claim 3, $uv^- \notin E(G)$ and $vu^- \notin E(G)$. Thus, y_1 and y_2 are well-defined.

Claim 4. Let $w \in \{w_1, \cdots, w_r\}, x \in \overrightarrow{C}[u^+, y_1]$ and $y \in \overrightarrow{C}[v^+, y_2]$. Then we have

(1) $wx \notin \widetilde{E}(G), wy \notin \widetilde{E}(G)$;

(2) $uy \notin \widetilde{E}(G), vx \notin \widetilde{E}(G)$;

(3) $xy \notin \widetilde{E}(G)$.

Proof. (1) Suppose that $wx \in \widetilde{E}(G)$. By Claim 1, $x \neq u^+$ and this implies that $ux^- \in E(G)$. Then $C' = \overrightarrow{P}[u, w]wx\overrightarrow{C}[x, u^-]u^-u^+\overrightarrow{C}[u^+, x^-]x^-u$ is an o-cycle longer than C and contains all vertices in C, a contradiction. The other assertion can be proved similarly.

(2) Suppose that $uy \in \widetilde{E}(G)$. By Claim 3, $y \neq v^+$ and this implies that $vy^- \in E(G)$. Then $C' = uy\overrightarrow{C}[y, u^-]u^-u^+\overrightarrow{C}[u^+, v^-]v^-v^+\overrightarrow{C}[v^+, y^-]y^-v\overleftarrow{P}[v, u]$ is an o-cycle longer than C and contains all vertices in C, a contradiction. Similarly, the other assertion can be proved.

(3) Suppose that $xy \in \widetilde{E}(G)$. By Claim 2, $u^-u^+ \in \widetilde{E}(G)$ amd $v^-v^+ \in \widetilde{E}(G)$. Now $C' = \overrightarrow{P}vy^-\overleftarrow{C}[y^-, v^+]v^+v^-\overleftarrow{C}[v^-, x]xy\overrightarrow{C}[y, u^-]u^-u^+\overrightarrow{C}[u^+, x^-]x^-u$ (if $x \neq u^+$ and $y \neq v^+$) or $C' = \overrightarrow{P}vy^-\overleftarrow{C}[y^-, v^+]v^+v^-\overleftarrow{C}[v^-, u^+]u^+y\overrightarrow{C}[y, u]$ (if $x = u^+$ and $y \neq v^+$) or $C' = \overrightarrow{P}\overleftarrow{C}[v, x]xv^+C[v^+, u^-]u^-u^+C[u^+, x^-]x^-u$ (if $x \neq u^+$ and $y = v^+$) is an o-cycle longer than C and contains all vertices in C, a contradiction. □

Claim 5. $u^-u^+ \in E(G)$ or $v^-v^+ \in E(G)$.

Proof. Suppose that $u^-u^+ \notin E(G)$ and $v^-v^+ \notin E(G)$. By Claim 2, we have $d(u^-) + d(u^+) \geq n$ and $d(v^-)+d(v^+) \geq n$. Thus, we obtain $d(u^-)+d(v^-) \geq n$ or $d(u^+)+d(v^+) \geq n$, contradicting to Claim 3. □

By Claim 5, without loss of generality, we assume that $u_{-1}u_1 \in E(G)$.

Case 1. $uv \notin E(G)$ or $uv \in E(G)$ and $r = 1$.

Suppose that $uv \notin E(G)$. By Claim 4, $\{y_1, y_1^-, u, w_1, \ldots, w_r, v, y_2^-, y_2\}$ induces a P_{6+r}, where $r \geq 1$. Since G is P_7-f-heavy, G is also P_{6+r}-f-heavy. By the choice of C, w_1 and w_r are light. It follows that y_1^- and y_2^- are heavy, and this implies $y_1^- y_2^- \in \widetilde{E}(G)$, contradicting to Claim 4 (3).

Suppose that $uv \in E(G)$ and $r = 1$. Recall that G is D-f-heavy or H-f-heavy. If G is D-f-heavy, then by Claim 4, $\{y_1, y_1^-, u, w_1, v, y_2^-, y_2\}$ induces a D. Since w_1 is light, y_1^- and y_2^- are heavy. It follows that $y_1^- y_2^- \in \widetilde{E}(G)$, which contradicts to Claim 4 (3). Now, we assume that G is H-f-heavy. By Claims 1 and 3, $\{u^-, u, u^+, w_1, v\}$ induces an H. It follows that u^- is heavy. If $v^- v^+ \in E(G)$, then by Claims 1 and 3, $\{v^-, v, v^+, w_1, u\}$ induces an H. Similarly, we have v^- is heavy. If $v^- v^+ \notin E(G)$, then $\{v^-, v, v^+, w_1\}$ induces a claw. Since w_1 is light and G is 2-heavy, we have v^- is heavy. In these two cases, we obtain $u^- v^- \in \widetilde{E}(G)$, which contradicts to Claim 3.

Case 2. $uv \in E(G)$ and $r \geq 2$.

Since $r \geq 2$ and the choice of the path P, we have $w_1 v \notin E(G)$ and $w_r u \notin E(G)$. By Claim 3, we obtain $uv^+ \notin E(G)$ and $u^- v \notin E(G)$. By Claim 1, $w_1 u^- \notin E(G)$ and $w_r v^+ \notin E(G)$. Thus each of $\{u, w_1, u^-, v\}$ and $\{v, w_r, v^-, u\}$ induces a claw. Since each of $\{w_1, w_r\}$ is light and G is 2-heavy, u^- and v^- are heavy. Hence $u^- v^- \in \widetilde{E}(G)$, which contradicts to Claim 3.

The proof is complete. $\square$

Acknowledgement

The author is supported by NSFC (No. 11271300) and the Doctorate Foundation of Northwestern Polytechnical University (cx201326).

References

[1] P. Bedrossian, G. Chen, R.H. Schelp, A generalization of Fan's condition for Hamiltonicity, pancyclicity, and Hamiltonian connectedness, *Discrete Math.* **115** (1993) 39-50.

[2] B. Bollobás and G. Brightwell, Cycles through specified vertices, *Combinatorica* **13** (1993) 147-155.

[3] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications, Macmillan London and Elsevier, New York (1976).

[4] H.J. Broersma, Z. Ryjáček and I. Schiermeyer, Dirac's minimum degree condition restricted to claws, *Discrete Math.* **167/168** (1997) 155-166.

[5] H.J. Broersma and H.J. Veldman, Restrictions on induced subgraphs ensuring hamiltonicity or pancyclicity of $K_{1,3}$-free graphs, in: Contemporary Methods in Graph Theory, BI Witt.-Verl., Mannheim-Wien-Zürich, (1990) 181-194.

[6] G. Chen, B. Wei and X. Zhang, Forbidden subgraphs and hamiltonian cycles, preprint, (1995).

[7] G. Chen, B. Wei and X. Zhang, Degree-light-free graphs and hamiltonian cycles, *Graph and Combin.* **17** (2001) 409-434.

[8] G. Fan, New sufficient conditions for cycles in graphs, *J. Combin. Theory Ser. B* **37** (1984) 221-227.

[9] R.J. Faudree, Z. Ryjáček and I. Schiermeyer, Forbidden subgraphs and cycle extendability, preprint, University of West Bohemia, (1994).

[10] B. Li, Z. Ryjáček, Y. Wang and S. Zhang, Pairs of heavy subgraphs for Hamiltonicity of 2-connected graphs, *SIAM J. Discrete Math.* **26 (3)** (2012) 1088-1103.

[11] G. Li, B. Wei and T. Gao, A structural method for hamiltonian graphs, *Australasian J. Combin.* **11** (1995) 257-262.

[12] B. Ning and S. Zhang, Ore- and Fan-type heavy subgraphs for Hamiltonicity of 2-connected graphs, arXiv: 1203.3915 [math.CO].

[13] R. Shi, 2-Neighborhoods and hamiltonian conditions, *J. Graph Theory* **16** (1992) 267-271.